arXiv:1108.3397v2 [quant-ph] 24 Oct 2011

Influence of measurement attributes on quantum correlations of the qubit pair with a common reservoir

A. Thilagam
Information Technology, Engineering and the Environment,
Mawson Institute, University of South Australia, Australia 5095

(Dated: October 25, 2011)

The effect of non-ideal or imperfect measuring attributes such as the quantum level of precision and finite duration on the classical correlation and quantum discord is analysed for a pair of qubits immersed in a common reservoir. We show that the quantum discord is enhanced as the precision of the measuring instrument is increased, and both the classical correlation and the quantum discord undergo noticeable changes during the duration when measurements are performed on a neighboring partition. These results are in stark contrast to those obtained using idealized measurement procedures, and imply the emergence of classicality in quantum systems with increasing imperfections in measurement procedures.

PACS numbers: 03.67.-a, 03.65.Ta, 03.65.Yz

I. INTRODUCTION

The role of quantum measurements in quantum entanglement, correlation and decoherence processes present one of the most challenging areas in the current context of quantum mechanical investigations in several systems [1–9]. Some progress has been made in the understanding of the links between quantum measurements and the decoherence process which results in the breakdown of phases in the superpositions of states that exist prior to measurement procedures [10]. In the "decoherence scheme," a quantum system which survives while in contact with an environment is guided to its pointer states [11]. The view that an open quantum system is equivalent to a system which is continuously measured by its environment has been pursued by other groups based on a variety of approaches [12, 13]. However, some questions still exist, for instance the quantum measurement stands distinct from other kinds of interactions in that it alone brings about the collapse of the wavefunction of the observed system, and the reasons for this remain unclear. A related problem is the detection of the exact point at which a classical world emerges from a group of interacting quantum systems, and the various factors which trigger such transitions. Recent works [14, 15] have shown that invasive measurements which have an influence on the dynamics of the monitored system can give rise to classicality in quantum systems.

Quantum measurements have taken more important roles in recent years due to their role in the definition of several measures of quantum correlations, including the quantum discord [16–18] which has been shown to possess more generalized properties than other well established measures such as the Wootters concurrence [19]. In order to evaluate the quantum discord, a set of positive-operator-valued measurements (POVM) is performed in a neighboring partition to determine the extent to which the system is changed by such measurements. At zero quantum discord, a measurement procedure allows external observers to obtain all information about a bipartite system without disturbing it. Two quantum systems can be considered as purely classically linked if the measurements performed on one system does not affect the adjacent system. To date, the quantum discord appears as a useful entity [20–24] in differentiating processes which are essentially quantum from those that incorporate generically non-classical features.

The quantum discord does not obey the well known monogamy rule that arises as a consequence of the linearity of quantum-mechanical systems. The quantum discord is defined conventionally based on orthogonal projections of ideal measurements [16–18]. It is not clear whether a non-ideal or invasive measurement that interferes with the dynamics of the measured system can distort the quantum discord. In this regard, there is a need to study the key attributes of the measurement process that can influence the quantum discord, and if necessary whether these imperfections can decrease the quantum correlations present in entangled systems. These considerations form the main motivation for this this work in which we introduce measurement attributes such as the duration as well as quantum level precision of the measurement procedure, and examine the effect of doing this on the resulting dynamics of a qubit pair immersed in a common reservoir. In contrast, ideal measurements are assumed to occur instantaneously with no intervention on the dynamics of the monitored system.

In order to keep the numerical analysis tractable, we adopt the Feynman's path integral framework [25, 26] to interpret quantum measurements, and in particular, employ a variant of this formalism based on the restricted path integral formalism[9, 27]. Within the restricted path framework, the continuous measurement of a quantity with a given result is monitored by constraints imposed on the Feynman's path integral, accordingly an anti-Hermitian term is added to the Hamiltonian describing the dynamics of the system during the measurement process. The restricted path integral is appealing as it is dependent solely on the information gained from the measurement procedure and is independent of the detailed construct of the measuring device. The dependence on parameters associated with a secondary system acting as the observer or measuring device is therefore reduced. Moreover, the results obtained is applicable to any quantum configuration of subsystems, which makes it convenient to analyse the critical dynamics associated with transfer of information between distinct quantum entities. As a consequence, these features give rise to simplicity in analytical treatment in situations when only the effect of a measurement needs to be incorporated within an open quantum system.

We recall the two key elements in Feynman's path integral formalism [25, 26], the first involves the superposition principle which yields the transition amplitude, and not the probability, for a given quantum process in the absence of measurements. Under this scheme, the probability amplitude of the transition from the initial to the final state of the system is obtained via summation of the amplitudes of all possible paths which could also interfere with each other. The second element in Feynman's formalism involves the weight attached to the individual paths which are involved in the summation, this weight provides a measure of contribution of each constituent path.

In open quantum systems, the degrees of freedom of the constituent paths undergo decoherence under the action of a measurement process, which disrupts the interference between the alternative paths by virtue of the uncertainty principle [28]. Unlike ideal measurements in which a quantity is measured without any disturbance to the system, weak measurements induce minimal disturbance of the various possible routes so that the interference between alternative paths remains almost intact [29, 30]. It is also likely that frequent measurement could not only hinder the dynamics of a system, but may also guide the evolution of the system. Hence, a sequence of measurements along a given path could induce the dynamics of the system to follow a selected trajectory [31, 32].

The Von Neumann projection operator $\mathcal{P}$ [4, 7] is an effective tool in formulating ideal measurement procedures in Hilbert space $\mathcal{H}$ of a quantum system. Within this scheme, the act of measurement decomposes the total Hilbert space into partitions of orthogonal quantum Zeno subspaces [7], and the initial density matrix ρ_0 is then confined to $\mathcal{H}_{\mathcal{P}}$ with a certain probability. Transitions outside this space is minimized for a large number of measurements N taken at regular time intervals. The survival probability becomes one at very large N as different outcomes are eliminated. In this regard, the Feynman path formalism and the notion of quantum Zeno subspaces [7] can be seen to lead to the same conclusion with regard to measurement and decoherence processes.

Highly precise measurements stand distinct from the less complex ideal or weak measurements in which observables manifests as Hermitian operators which act on the state space to yield the eigenvalue of the eigenstate at which the system existed during the period of measurement. In highly precise measurements, the quantum evolution follows a complex route due to the influence of non-Hermitian terms, which interfere strongly with the dynamics of the measured system, and may even dominate the time evolution of the quantum system. Hence, it is not immediately clear what reading will be obtained and at which state the monitored system is actually existing in after a highly precise measurement. Moreover, it is implicit that orthogonal measurement projections may themselves be in a state of evolution during a measurement process which is of finite duration. To investigate some of these issues, we include non-Hermitian terms in in this work, and examine how the quantum discord is affected if monitoring device imparts a significant disturbance on the system under study, and itself dominates the time evolution of the quantum system. The key feature of this work is therefore, the possibility that projections can be still made on imperfect measurements. This procedure which departs from conventional treatments, is not illogical in view of the statistical nature of quantum processes.

This paper is organized as follows. In Section II we describe the restricted path integral approach for energy measurements, which incorporates the non-ideal attributes of the measuring device. These non-ideal features of the measurement device lead to interference of the dynamics of the quantum system under observation, hence the effect of non-ideal measurement in entangled states is also discussed in this section. In Section III, the classical correlation and quantum discord measures are discussed, and a brief description of the qubit pair system immersed in a common reservoir is described. In Section IV, the effect of the measurement precision and finite time duration on the quantum correlation measure is evaluated and numerical results are presented as well. A brief discussion and the conclusion are also presented in this Section.

II. THE RESTRICTED PATH INTEGRAL PATH APPROACH FOR ENERGY MEASUREMENTS

The restricted path integral is derived [9, 27, 33, 34] from the Feynman path integral via the introduction of a weight functional within the integrand of the various paths involved in the summation process. We recall that the Feynman's propagator, $K_{[E]}(q', \tau; q, 0)$ in the phase-space representation at time τ is given by [25, 26]

$$K(q', \tau; q, 0) = \int d[q]d[p]e^{\frac{i}{\hbar}\int_0^\tau [p\dot{q} - \widehat{H}_0(q,p)]dt} \tag{1}$$

where $\widehat{H}_0$ is the Hamiltonian of the closed (unmeasured) quantum system and $[p]$ and $[q]$ are the paths in the momentum and configuration spaces respectively. In Mensky's formalism, the output of a quantum system subjected to measurement is expressed in terms of constrained paths linked to the measured system via a weight functional $w_{[E]}$ [9]. The functional may assume a Gaussian form with a damping magnitude that is proportional to the squared difference between the observed value along the paths and the actual measurement result. Thus a system subjected to measurement evolves via a propagator which modifies Eq.(1) according to [34, 35]

$$K_{[E]}(q',\tau;q,0)=\int d[q]d[p]e^{\frac{i}{\hbar}\int_0^\tau[p\dot{q}-\widehat{H}_0(q,p)]dt}w_{[E]} \tag{2}$$

Eq. (2) is dependent on a selected measurement output such as E after a time τ, for a measuring instrument that incurs an error E_r during the measurement.

Using Eq.(2), it can be shown [35–38] that the stimulated transition in a two-level system is affected by the sensitivity or error in measurements of its energy levels by an external device. These results show that a highly precise measurement will result in a stagnation of transitions in the two-level system, which is partially consistent with the quantum zeno effect[4, 6, 7]. In a recent work [38], singularities known as exceptional points [39] are shown to appear at the branch point of eigenfunctions at a critical measurement precision E_r^c. E_r^c is determined by parameters of the two-level system and measurement duration τ [38]. The significance of the results from Mensky's formalism lies in the inclusion of attributes of the measuring device, which determines the dynamics of the quantum system under observation. This has obvious implications for the evaluation of the quantum discord in quantum systems, as will be shown later in this work.

The use of the Gaussian measure, $w_{[E]}=\exp\left\{-\frac{\langle(H_0-E)^2\rangle}{\Delta E^2}\right\}$ enables the effect of the measurement to be incorporated via the effective Hamiltonian [35, 36] for a two-level system

$$\widehat{H}_{eff}=\widehat{H}_0-i\frac{\hbar}{\tau E_r^2}(\widehat{H}_0-E)^2 \tag{3}$$

where $\langle ...\rangle$ denotes the time-average for the duration τ during which measurement was performed. As noted earlier, E is the selected measurement output after a time τ and E_r is the error made during the measurement of the energy, E. E_r can therefore be considered as a measure of the precision of the monitoring device. A large (small) error made during the measurement can be viewed as a weak (highly precise) measurement. The system therefore evolves as $i\hbar\frac{\partial}{\partial t}|\psi(t)\rangle=H_{eff}|\psi(t)\rangle$ during measurement. By expanding the state of the system being measured within the unperturbed basis states $|n\rangle$ of the unmeasured system with Hamiltonian $\widehat{H}_0$ as $|\psi(t)\rangle=\sum_n C_n(t)|n\rangle$, the coefficients $C_n(t)$ can be solved via the Schrödinger equation Eq.(3).

A. Effect of non-ideal measurement in entangled states

We now consider the application of the restricted path approach to the entangled states in a two-level system with energies E_1 (E_2) at state $|0\rangle$ ($|1\rangle$) and subjected to a continuous measurement process. The two-level qubit Hamiltonian in the ideal unmeasured scenario is written as

$$\hat{H}_0=-\hbar(\frac{\Delta\omega}{2}\,\sigma_z+V(t)\,\sigma_x), \tag{4}$$

where the Pauli matrices $\sigma_x=|0\rangle\langle 1|+|1\rangle\langle 0|$ and $\sigma_z=|1\rangle\langle 1|-|0\rangle\langle 0|$. The external perturbation $V(t)$ incorporates the transitions between the two levels and $\Delta\omega=2(E_1+E_2)$. The perturbation potential $V_{00}=V_{11}=0$ and $V_{01}=V_{10}^*=V_0e^{i\omega(t-t_0)}$ with V_0 real. For the two-level system subjected to a measurement process, the first term in Eq. (4) transforms via the non-Hermitian Hamiltonian in Eq.(3) due to the constraining effect of the selected readout E. In the presence of the external perturbation, the total Hamiltonian in Eq.(4) also evolves via the effective Hamiltonian in Eq.(3), but with inclusion of the external perturbative term $V(t)$ as follows

$$|\psi(t)\rangle=e^{-i(E_1-2i\lambda_1)t}C_1(t)\,|0\rangle+e^{-i(E_2-2i\lambda_2)t}C_2(t)\,|1\rangle \tag{5}$$

where $\lambda_1=\frac{(E_1-E)^2}{2\tau E_r^2}$ and $\lambda_2=\frac{(E_2-E)^2}{2\tau E_r^2}$.

The coefficients $C_1(t),C_2(t)$ in Eq.(5) can be obtained as

$$\begin{bmatrix} C_1(t) \\ C_2(t) \end{bmatrix}=\begin{bmatrix} \cos\kappa t-i\alpha_1 & -i\alpha_2 \\ -i\alpha_2 & \cos\kappa t+i\alpha_1 \end{bmatrix}\begin{bmatrix} C_1(0) \\ C_2(0) \end{bmatrix}, \tag{6}$$

where α_1=$\cos\theta\sin\kappa t$, α_2=$\sin\theta\sin\kappa t$, $\cos\theta$=$\frac{q}{\kappa}$, κ=$\sqrt{q^2+V_0^2}$, q=$\frac{1}{2}(\omega-\Delta E+2i\Omega)$, ΔE=(E_2-E_1), and Ω=λ_2-λ_1. The terms λ_2 and λ_1 as defined in the earlier paragraph are dependent on the measurement precision, E_r as well as the energy E (E_1 or E_2) to be measured.

The entangled states of the monitored system therefore incorporate non-Hermitian terms[38, 40, 41]

$$\begin{aligned} |\chi_{\rm s}(t)\rangle &= e^{-\lambda_r t/2}\left(\cos\kappa t - i\cos\theta\sin\kappa t\right)|0\rangle \\ &\quad -ie^{-\lambda_r t/2}\sin\theta\sin\kappa t\,|1\rangle \\ |\chi_{\rm a}(t)\rangle &= e^{-\lambda_r t/2}\left(\cos\kappa t + i\cos\theta\sin\kappa t\right)|1\rangle \\ &\quad -ie^{-\lambda_r t/2}\sin\theta\sin\kappa t\,|0\rangle\,, \end{aligned} \tag{7}$$

where λ_r=$\frac{\Delta E^2}{2\tau E_r^2}$. For measurement procedures which introduce very large errors, $E_r\to\infty$,λ_1=λ_2=λ_r=$\cos\theta$=0, and the excitonic qubit oscillates coherently between the two levels with the Rabi frequency $2V_0$ as is well known in undisturbed systems.

We consider the system in which the initial state at $t=0$ is $|1\rangle$ and the final state at time t is either $|1\rangle$ or $|0\rangle$, the probabilities P_{11} (P_{10}) of the system to be in the state $|1\rangle$ ($|0\rangle$) can be obtained depending on the absolute values of V_0 and λ_r=$\frac{(E_2-E_1)^2}{2\tau E_r^2}$ [38, 41] . At the resonance frequencies, $\omega=\Delta E$, κ=κ_0 and there are two regimes, depending on the relation between V_0 and λ_r. The range where $V_0>\lambda_r$ applies to the coherent tunneling regime and we obtain $P_{11}=e^{-\lambda_r t}[\cos\kappa_0 t-\frac{\lambda_r}{\kappa_0}\sin\kappa_0 t]^2$, $P_{10}=e^{-\lambda_r t}\frac{V_0^2}{\kappa_0^2}\sin^2\kappa_0 t$. For $V_0<\lambda_r$, the system undergoes incoherent tunneling and $P_{11}=e^{-\lambda_r t}[\cosh\kappa t-\frac{\lambda_r}{\Omega_0}\sinh\kappa t]^2$, $P_{10}=e^{-\lambda_r t}\frac{V_0^2}{\kappa_0^2}\sinh^2\kappa_0 t$. At the singularities known as exceptional points, $\kappa_0=0$, and both regimes merge and we obtain $P_{11}=\left(1-\frac{\lambda_r t}{2}\right)^2 e^{-\lambda_r t}$ and $P_{10}=\left(\frac{\lambda_r t}{2}\right)^2 e^{-\lambda_r t}$ [39]. The exceptional point appears at a critical measurement precision E_r^c=$\frac{\Delta E}{\sqrt{2\tau V_0}}$. Considering a unit system in which $\hbar=V_0=\Delta E$=1, $\tau=2\pi/V_0$ and a unitless time $t=t'/\tau$, we obtain $E_r^c=\frac{1}{\sqrt{4\pi}}$. Using r to denote the unitless measurement precision parameter, we note that at $r>\frac{1}{\sqrt{4\pi}}$ ($r<\frac{1}{\sqrt{4\pi}}$), the quantum system undergoes coherent (incoherent) tunneling.

III. CLASSICAL CORRELATION AND QUANTUM DISCORD

Following the formulation of quantum discord in Refs.[16–18], we express the quantum mutual information of a composite state ρ of two subsystems A and B as $\mathcal{I}(\rho)=S(\rho_A)+S(\rho_B)-S(\rho)$ for a density operator in $\mathcal{H}_A\otimes\mathcal{H}_B$. ρ_A (ρ_B) is the reduced density matrix associated with A (B) and $S(\rho_i)$ (i=A,B) denotes the well known von Neumann entropy of the density operator ρ_i, where $S(\rho)=-\mathrm{tr}(\rho\log\rho)$. The mutual information can also be written in terms of quantum conditional entropy $S(\rho|\rho_A)=S(\rho)-S(\rho_A)$ as

$$\mathcal{I}(\rho)=S(\rho_B)-S(\rho|\rho_A) \tag{8}$$

A series of one-dimensional orthogonal projectors $\{\Pi_k\}$ induced in $\mathcal{H}_A$ leads to different outcomes of the measurement in $\mathcal{H}_B$ via the post measurement conditional state

$$\rho_{B|k}=\frac{1}{p_k}(\Pi_k\otimes\mathbb{I}_B)\rho(\Pi_k\otimes\mathbb{I}_B) \tag{9}$$

where the probability $p_k=\mathrm{tr}[\rho(\Pi_k\otimes\mathbb{I}_B)]$ and $\{\Pi_k\}$ denote the one-dimensional projector indexed by the outcome k. From the cumulative effect of the mutually exclusive measurements on A, we obtain a conditional entropy of the subsystem B based on $\rho_{B|k}$

$$S(\rho|\{\Pi_k\})=\sum_k p_k S(\rho_{B|k}) \tag{10}$$

which is used to obtain the measurement induced mutual information $\mathcal{I}(\rho|\{\Pi_k\})=S(\rho_B)-S(\rho|\{\Pi_k\})$. The classical correlation measure based on optimal measurements made on A is obtained as [16–18]

$$\mathcal{C}_A(\rho)=\sup_{\{\Pi_k\}}\mathcal{I}(\rho|\{\Pi_k\}) \tag{11}$$

The difference in $\mathcal{I}(\rho)$ and $\mathcal{C}_A(\rho)$ yields the non symmetric quantum discord $\mathcal{D}_A(\rho)=\mathcal{I}(\rho)-\mathcal{C}_A(\rho)$ A discord $\mathcal{D}_B(\rho)$ corresponding to measurements made on B can likewise be obtained and need not be the same as $\mathcal{D}_A(\rho)$. We note that measurements made on a neighboring partition is the key to determining the classical correlation measure between the subsystems.

A. A qubit pair immersed in a common reservoir

The joint evolution of a pair of two-level qubit subsystems, A, B undergoing decoherence in a common reservoir is determined by a completely positive trace preserving map expressed in the operator-sum form [42]

$$\varepsilon\left(\rho_{AB}\right) = \sum_{i,j} \Gamma_i(A)\Gamma_j(B)\; \rho_{AB}\; \Gamma_i^\dagger(B)\Gamma_j^\dagger(A), \tag{12}$$

where $\Gamma_i(A)$ ($\Gamma_i(B)$) is the Kraus operator associated with the decoherence process at A (B). For the phase flip channel in which there is loss of quantum information with conservation of energy, the Kraus operators are given in the basis $\{\left|0\right\rangle, \left|1\right\rangle\}$ for both subsystems, $k = A, B$ as [21, 42] $\Gamma_0(A) = diag(\sqrt{1-p/2}, \sqrt{1-p/2}) \otimes \mathbf{1}_B$, $\Gamma_1(A) = diag(\sqrt{p/2}, -\sqrt{p/2}) \otimes \mathbf{1}_B$, $\Gamma_0(B) = \mathbf{1}_A \otimes diag(\sqrt{1-p/2}, \sqrt{1-p/2})$ and $\Gamma_1(B) = \mathbf{1}_A \otimes diag(\sqrt{p/2}, -\sqrt{p/2})$, The parameter $p = 1 - \exp(-\gamma t)$, where γ denotes the phase damping rate.

To simplify the numerical analysis, we consider the joint state of a pair of two-level qubit subsystems, A, B to be prepared in the initial X-type state with maximally mixed marginals ($\rho_{A(B)} = I_{A(B)}/2, S(\rho_A(t)) = S(\rho_B(t)) = 1$) with density matrix of the form $\rho(0) = \frac{1}{4}[I + \sum\limits_{i=1}^{3} c_i \sigma_A^i \otimes \sigma_B^i]$. where I is the identity operator associated with the qubit pair, σ_j^i and σ ($j = A, B$) are the Pauli operators of each qubit. c_i ($0 \leq |c_i| \leq 1$) are real numbers, with the Werner states sharing a common $|c_1| = |c_2| = |c_3| = c$ and c=1 for the Bell basis states. We assume that the usual unit trace and positivity conditions of the density operator ρ are satisfied. For the class of states where $|c_1| = |c_2| = c, |c_3| = c_3$), the evolution of the joint system is described by the matrix

$$\rho_{\text{A,B}}(t) = \frac{1}{4} \begin{pmatrix} 1+c_3 & 0 & 0 & 0 \\ 0 & 1-c_3 & 2ce^{\mu^* t} & 0 \\ 0 & 2ce^{\mu t} & 1-c_3 & 0 \\ 0 & 0 & 0 & 1+c_3 \end{pmatrix}, \tag{13}$$

where $\mu = \left[-2\gamma - i(\Delta\omega_A - \Delta\omega_B)\right] t$. As noted earlier, which γ is the phase damping rate. To simplify the analysis, we have considered the same damping rate in both qubit subsystems. $\Delta\omega_i, i$ =A,B denotes the difference in energy levels of each qubit subsystem, we consider equivalent energy levels in the qubit pair in this work. The mutual information of state $\rho_{\text{A,B}}$ in Eq. (13) is evaluated using $\mathcal{I}\left(\rho_A : \rho_B\right) = 2 + \sum_{i=1}^{4} \lambda_i \log \lambda_i$ where the eigenvalues λ_i of $\rho_{\text{A,B}}$ are $\lambda_{1,2} = \frac{1}{4}(1+c_3), \lambda_3 = \frac{1}{4}(1 - c_3 + 2ce^{-2\gamma t})$ and $\lambda_4 = \frac{1}{4}(1 - c_3 - 2ce^{-2\gamma t})$

IV. MEASUREMENT PRECISION AND QUANTUM CORRELATIONS

The classical correlation measure is obtained through all possible local measurements on one of the subsystems, say A. For the ideal case of a local measurement that is instantaneous, we utilize a set of orthogonal projectors $\{\Pi_k = \left|\theta_k\right\rangle \left\langle\theta_k\right| |, k = \parallel, \perp\}$, which are defined in terms of the orthogonal states

$$\begin{aligned} \left|\theta_\parallel\right\rangle &= \cos\theta \left|0\right\rangle + e^{i\phi} \sin\theta \left|1\right\rangle, \\ \left|\theta_\perp\right\rangle &= e^{-i\phi} \sin\theta \left|0\right\rangle - \cos\theta \left|1\right\rangle, \end{aligned} \tag{14}$$

where $0 \leq \theta \leq \pi/2$ and $0 \leq \phi \leq 2\pi$. In a recent work, Galve et. al. [43] showed that orthogonal measurements are sufficient to evaluate the quantum discord pertaining to rank 2 states of two qubit systems, but provide tight upper bounds for higher rank (3 and 4) states.

In the case involving imperfect measurements of finite duration, τ and precision E_r, we consider that measurements can still be modeled via projector operations. This implies that the orthogonal measurement projections may themselves be in a state of evolution during the measurement process. Using Eq. (7) which highlight the effect of non-ideal measurements on a single qubit as basis, we transform the orthogonal states in Eq. (14) as

$$\begin{aligned} \left|\theta_\parallel\right\rangle_p &= R(\theta) \left|0\right\rangle + e^{i\phi} S(\theta) \left|1\right\rangle, \\ \left|\theta_\perp\right\rangle_p &= e^{-i\phi} S(\theta) \left|0\right\rangle - R(\theta) \left|1\right\rangle, \end{aligned} \tag{15}$$

where the terms $R(\theta) = e^{-\lambda_r t/2}\left(\cos\theta t - i\frac{\lambda_r t}{\theta}\sin\theta\right)$ and $S(\theta) = e^{-\lambda_r t/2}\frac{\sqrt{\theta^2 + (\lambda_r t)^2}}{\theta}\sin\theta$. As noted earlier, $\lambda_r = \frac{\Delta E^2}{2\tau E_r^2}$ and ΔE is the energy difference between the $\left|1\right\rangle$ and $\left|0\right\rangle$ states of the qubit. In the limit of $\lambda_r \to 0$, Eq. (15) reduce to the orthogonal set in Eq. (14). The dependence of the projected states given in Eq. (15) on the measurement attributes, τ and λ_r results in the dependence of the classical correlation on these same attributes, and we examine such a relationship next.

The eigenvalues of the reduced density matrix, $\rho_B^{(k)}$ of the neighboring subsystem B in accordance with the two projective measurements ($p_{\shortparallel} = p_\perp = 1/2$) in subsystem A are obtained using Eqs. (15) as

$$\begin{aligned} \zeta_{1,2}^{(k)} &= \frac{1}{2}(1 \pm \Theta), \qquad (16) \\ \Theta^2 &= 4c_3^2 e^{-2pt}\left[\cos^2\theta + \frac{2pt\xi}{\theta^2}\sin^2\theta\right]^2 \\ &+ \frac{\xi^2}{\theta^2}c^2 e^{-4gt}e^{-2pt}\left[\cos^2\theta + \frac{(pt)^2}{\theta^2}\sin^2\theta\right]\sin^2\theta \end{aligned}$$

where $g = \gamma\tau$, $\xi = \sqrt{\theta^2 + (pt)^2}$, the unitless time, t is obtained via division with $\tau = 2\pi/V_0$ and the dimensionless precision parameter, p=$\lambda_r\tau$. Using Eq. (16), we obtain $S(\rho_B^{\|}) = S(\rho_B^{\perp}) = -\mathrm{tr}(\rho\log\rho) = R(\Theta) = -\frac{1-\Theta}{2}\log_2\left[\frac{1-\Theta}{2}\right] - \frac{1+\Theta}{2}\log_2\left[\frac{1+\Theta}{2}\right]$. The classical correlation given in Eq. (11) is evaluated using

$$\mathcal{C}(\rho) = 1 - \min_{\theta,\phi} R(\Theta), \qquad (17)$$

Further evaluation of $\mathcal{C}(\rho)$ is simplified by the elimination of the parameter ϕ as a result of the choice, $|c_1| = |c_2| = c$ used earlier for simplicity in analysis. The maximal value of Θ is thus not only dependent on c_3, c, γ, but also on the precision parameter p and the measurement duration τ. It is obvious from Eq. (16), that the influence of γ, the phase damping rate becomes critical only for $c > c_3$.

The quantum discord, which will also be influenced by the measurement attributes is given by

$$\mathcal{D}(\rho) = 2 + \sum_{k=1}^{4} \lambda_k \log_2 \lambda_k - \mathcal{C}(\rho), \qquad (18)$$



FIG. 1: a) Classical correlation $\mathcal{C}$ as function of t ($0 < t < \tau$) for various values of the measurement precision p. τ the measurement time duration is set at 1 and $g = \gamma\tau$=0.6 where γ is the phase damping rate. The real numbers $c_1 = c_2 = c$=0.15 and c_3=0.6. The curves from top to bottom correspond to the unitless measurement precision, p=0, 0.05, 0.2, 0.5.
b) Quantum discord $\mathcal{D}$ as function of t ($0 < t < \tau$) for various values of the measurement precision p=0.5, 0.2, 0.05, 0 (top to bottom). All other parameters are the same as in (a).

Figure 1a shows the changes in the classical correlation $\mathcal{C}$, obtained via numerically evaluation using Eqs. (16) and (17), for time range $0 < t < \tau$ for various values of the measurement precision p. Figure 1b illustrates changes in the quantum discord $\mathcal{D}$ which is evaluated using Eqs. (18) and the eigenvalues λ_i of $\rho_{\mathrm{A,B}}$ in Eq. (13). The figures shows that both $\mathcal{C}$ and $\mathcal{D}$ undergo noticeable changes during the duration when measurements are performed on a neighboring partition at non-zero p. Figure 1b shows that the quantum discord $\mathcal{D}$ is enhanced due to a corresponding decrease in $\mathcal{C}$ as p is increased. This trend is also reflected in Figure 2a,b which illustrates the changes in $\mathcal{C}$ and $\mathcal{D}$ with respect to the dephasing rate g for various values of the measurement precision p. We note in Figure 2a,b that beyond a critical g, the classical correlation $\mathcal{C}$ becomes independent of g as also noted in earlier works [21, 22] in which p=0.

In conclusion, we have presented results of the influence of non-ideal attributes such as the measurement precision and finite measurement time duration on the classical correlation and quantum discord for a qubit pair immersed



FIG. 2: a) Classical correlation $\mathcal{C}$ as function of the unitless $g = \gamma\tau$ where γ is the phase damping rate, for various values of the measurement precision p at t=1. The real numbers $c_1 = c_2 = c$=0.4 and c_3=0.2. The curves from top to bottom correspond to the unitless measurement precision, p=0, 0.2, 0.7.
b) Quantum discord $\mathcal{D}$ as function of g at t=1. for various values of the measurement precision p=0.7, 0.2, 0 (top to bottom). All other parameters are the same as in (a).

in a common environment. The results obtained show that the quantum discord is enhanced as the precision of the measuring instrument is increased, and both the classical correlation and the quantum discord undergo noticeable changes during the duration when measurements are performed on a neighboring partition. This work contributes to ideas proposed in recent works [14, 15] which showed the emergence of classicality from the quantum world for imprecise measurements. It is likely that in quantum measurements, which interfere with the dynamics of the measured system, the classical features from which classicality arise is enhanced. Moreover, the classicality may be linked with the breakdown in the number of alternative Feynman's routes from the initial to the final state of the system, however these ideas need further scrutiny and investigation. Lastly, the outcome of this study may provide useful guidelines in experimental techniques and studies of quantum measurement in optics and nanostructure systems [44, 45] and in experimental investigations involving non-classical correlations and emergence of classicality in entangled systems.

Acknowledgments

I would like to thank members of the MMaST group, University of South Australia, for their hospitality while this work was being carried out. This research was undertaken on the NCI National Facility in Canberra, Australia, which is supported by the Australian Commonwealth Government.

---

[1] J. von Neumann, "Mathematical Foundations of Quantum Mechanics," Princeton University Press, Princeton, (1955).
[2] H.D.Zeh, Found. Phys. **1**, 69 (1970).
[3] A. G. Kofman, G. Kurizki, Nature **405**, 546 (2000); Phys. Rev.A **54**, 3750(R)(1996).
[4] B. Misra and E. C. G. Sudarchan, J. Math. Phys. **18**, 758 (1977).
[5] P. Busch, P. J. Lahti, and P. Mittelstaedt, "The Quantum Theory of Measurement," Springer-Verlag, Berlin, (1991).
[6] W.M. Itano, D. J. Heinzen, J. J. Bollinger and D. J. Wineland, Phys. Rev.A **41**, 2295 (1990).
[7] P. Facchi and S. Pascazio, J. Phys. A: Math. Theor. **41** 493001 (2008); P. Facchi and S. Pascazio, Phys. Rev. Lett. **89**, 080401 (2002).
[8] V. B. Braginsky and F. Ya. Khalili, "Quantum Measurement", K. S. Thorne editor (Cambridge University Press, Cambridge) (1992), and references cited therein.
[9] M. B. Mensky, "Continuous Quantum Measurements and Path-Integrals" (Institute of Physics Publishers, Bristol and Philadelphia) (1993).
[10] W. H. Zurek, Phys. Today **44 (10)**, 36 (1991).
[11] W. H. Zurek, Phys. Rev. D **26**, 1862 (1982).
[12] M. Schlosshauer, "Decoherence and the Quantum-to-Classical Transition", Springer-Verlag, (2008).
[13] R. Ruskov and A. N. Korotkov, Phys. Rev. B **67**, 241305(R) (2003).
[14] J. Kofler and C. Brukner, Phys. Rev. Lett. **99**, 180403 (2007).
[15] J. Kofler and C. Brukner, Phys. Rev. Lett. **101**, 090403 (2008).
[16] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
[17] L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001).
[18] V. Vedral, Phys. Rev. Lett. **90**, 050401 (2003).

[19] V. Coffman, J. Kundu and W. K. Wootters, Phys. Rev. A **61**, 052306 (2000).
[20] A. Brodutch and D.R. Terno, Phys. Rev. A **81**, 062103 (2010).
[21] J. Maziero, L. C. Céleri, R. M. Sera and V. Vedral, Phys.Rev. A **80**, 044102 (2009).
[22] L. Mazzola, J. Piilo and S. Maniscalco, Phys. Rev. Lett. **104**, 200401 (2010).
[23] L. Ciliberti, R. Rossignoli and N. Canosa, Phys. Rev. A **82**, 042316 (2010).
[24] A. Datta, A. Shaji, and C. M. Caves, Phys. Rev. Lett. **100**, 050502 (2008).
[25] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, (McGraw-Hill, New York) (1965).
[26] R. P. Feynman, Rev. Mod. Phys. **20**, 367 (1948).
[27] M. B. Mensky, Phys. Rev. D **20**, 384 (1979); Sov. Phys. JETP **50**, 667 (1979).
[28] D. Sokolovski, I. Puerto Giménez and R. Sala Mayato, Phys.Lett.A **372**, 3784 (2008).
[29] Y. Aharonov, D. Z. Albert and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988).
[30] Y. Aharonov and L. Vaidman, J. Phys. A **24**, 2315 (1991).
[31] Y. Aharonov and M. Vardi, Phys. Rev. D **21**, 2235 (1980).
[32] T. P. Altenmuller and A. Schenzle, Phys. Rev. A **48**, 70 (1993).
[33] M. B. Mensky, R. Onofrio, and C. Presilla, Phys. Rev. Lett. **70**, 2825 (1993).
[34] M. B. Mensky, R. Onofrio, and C. Presilla, Phys. Lett. A **161**, 236 (1991).
[35] R. Onofrio, C. Presilla, and U. Tambini, Phys. Lett. A **183**, 135 (1993).
[36] U. Tambini, C. Presilla, and R. Onofrio, Phys. Rev. A **51**, 967 (1995).
[37] J. Ausretsch and M. B. Mensky, Phys. Rev. A **56**, 44 (1997).
[38] A. Thilagam, “Quantum Zeno effect on Quantum Discord”, arXiv:1011.5901v4 (2011)
[39] W. D. Heiss, Phys. Rev. E **61**, 929 (2000).
[40] J. C. Garrison and E. M. Wright, Phys. Lett. A **128**, 177 (1988).
[41] C. A. Stafford and B. R. Barrett, Phys. Rev. C **60**, 051305 (1999).
[42] A. Salles, F. de Melo, M. P. Almeida, M. Hor-Meyll, S. P. Walborn, P. H. Souto Ribeiro, and L. Davidovich, Phys. Rev. A **78**, 022322 (2008).
[43] F. Galve, G. L. Georgi and R. Zambrini, “Orthogonal measurements are almost sufficient for quantum discord of two qubits”, arXiv:1107.2005 v2 (2011)
[44] M. A. Hall, J. B. Altepeter and P. Kumar, Optics Express **17**, 14558 (2009).
[45] W. Tittel, J. Brendel, H. Zbinden, and N. Gisin, Phys. Rev. Lett. **81**, 3563 (1998).